

Review Practice: Chapter 14

Formulas:

$$z - z_0 = f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0)$$

$$L(x, y) \approx f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b)$$

$$\mathbf{D}_{\mathbf{u}}f(x, y, z) = \nabla f(x, y, z) \cdot \mathbf{u}$$

$$\text{Second Derivative Test: } D = \begin{vmatrix} f_{xx} & f_{xy} \\ f_{yx} & f_{yy} \end{vmatrix} = f_{xx}f_{yy} - (f_{xy})^2$$

$$\text{Lagrange Multipliers: } \nabla f(x, y, z) = \lambda \nabla g(x_0, y_0, z_0), \quad g(x, y, z) = k$$

$$\frac{\partial y}{\partial x} = -\frac{F_x}{F_y}, \quad \frac{\partial z}{\partial x} = -\frac{F_x}{F_z}, \quad \frac{\partial z}{\partial y} = -\frac{F_y}{F_z}$$

$$F_x(x_0, y_0, z_0)(x - x_0) + F_y(x_0, y_0, z_0)(y - y_0) + F_z(x_0, y_0, z_0)(z - z_0) = 0$$

- True or False: If $(2, 1)$ is a critical point of f and $f_{xx}(2, 1)f_{yy}(2, 1) < f_{xy}(2, 1)^2$ then f has a saddle point at $(2, 1)$.
- Find the partial derivatives of $G(x, y, z) = e^{xy} \sin(y/z)$.

3. If $v = x^2 \sin y + ye^{xy}$ where $x = s + 2t$ and $y = st$, use the chain rule to find $\frac{\partial v}{\partial t}$ and $\frac{\partial v}{\partial s}$ when $s = 0$ and $t = 1$.
4. Find the maximum rate of change of $f(x, y) = x^2y + \sqrt{x}$ at the point $(2, 1)$. In what direction does it occur?
5. Use Lagrange Multipliers to find the max/min values of $f(x, y) = \frac{1}{x} + \frac{1}{y}$ subject to the constraint $\frac{1}{x^2} + \frac{1}{y^2} = 1$. [Note I will only ask you for the set up of these problems]

6. Find the absolute max and min values of f on the set D given:

$$f(x, y) = e^{-x^2-y^2}(x^2 + 2y^2) \quad \text{and} \quad D = \{(x, y) \mid x^2 + y^2 \leq 4\}$$