

Review Practice: Chapters 14 & 15

1. Consider $f(x, y) = x^2 + 2x - y$

(a) Find all first and second partials

$$f_x = 2x + 2, \quad f_y = -1; \quad f_{xx} = 2, \quad f_{yy} = 0, \quad f_{xy} = f_{yx} = 0$$

(b) Find the gradient

$$\nabla f = \langle 2x + 2, -1 \rangle$$

(c) What types of graphs are the level curves of f ?

$$K = x^2 + 2x - y$$

$$y = x^2 + 2x + K \rightarrow \text{upward parabolas}$$

$$y = (x + 1)^2 + (K - 1) \quad \text{vertex: } (-1, K - 1)$$

2. Find all critical points of $f(x, y) = x^3 - 12x + y^2$ and classify them using the second derivative test.

$$\vec{0} = \nabla f = \langle 3x^2 - 12, 2y \rangle$$

$$x = \pm 2, \quad y = 0$$

$$D = f_{xx} \cdot f_{yy} - f_{xy}^2 = (6x)(2) - 0 = 12x$$

• $D(2, 0) > 0, \quad f_{xx}(2, 0) > 0 \Rightarrow$ local min at $(2, 0)$


• $D(-2, 0) < 0 \Rightarrow$ Saddle point at $(-2, 0)$

3. If $x = f(r, s)$ and $y = h(r, s)$ use chain rule to find $\frac{\partial z}{\partial s}$

$$\frac{\partial z}{\partial s} = f_x \frac{\partial x}{\partial s} + f_y \frac{\partial y}{\partial s}$$

$$\frac{\partial z}{\partial s} = f_x(g, h)g_s + f_y(g, h)h_s$$

4. Compute by changing the order of integration: $\int_0^1 \int_x^1 y^2 \sin(xy) \, dy \, dx$



$$= \int_0^1 \int_0^y y^2 \sin(xy) \, dx \, dy$$


$$= \int_0^1 -y^2 \frac{\cos(xy)}{y} \Big|_0^y \, dy$$

$$= \int_0^1 -y \cos(y^2) + y \, dy$$

$$= -\frac{\sin(y^2)}{2} + \frac{y^2}{2} \Big|_0^1 = \frac{1}{2} (-\sin(1) + 1)$$


5. Compute $\int \int_D x \, dA$ where D is the region in the first quadrant between $x^2 + y^2 = 1$ and $x^2 + y^2 = 2$.

$$\begin{aligned} \iint_D x \, dA &= \int_0^{\pi/2} \int_1^{\sqrt{2}} r \cos \theta \cdot r \, dr \, d\theta = \int_0^{\pi/2} \cos \theta \, d\theta \cdot \int_1^{\sqrt{2}} r^2 \, dr \\ &= \sin \theta \Big|_0^{\pi/2} \cdot \frac{r^3}{3} \Big|_1^{\sqrt{2}} \\ &= \frac{1}{3} \left(2^{3/2} - 1 \right) \end{aligned}$$



6. Compute $\iiint_E z \, dV$ where E is the region in the first octant between $y^2 + z^2 = 1$ and $x + y = 2$.

$$\begin{aligned} \iiint_E z \, dV &= \iint_D \int_0^{2-y} z \, dx \, dA \\ &= \iint_D (2z - yz) \, dA = \int_0^{\pi/2} \int_0^1 (2r \sin \theta - r^2 \sin \theta \cos \theta) \, dr \, d\theta \\ &= \int_0^{\pi/2} \left(\frac{2}{3} \sin \theta - \frac{1}{4} \sin \theta \cos \theta \right) \, d\theta = -\frac{2}{3} \cos \theta - \frac{1}{4} \frac{\sin^2 \theta}{2} \Big|_0^{\pi/2} = \boxed{-\frac{1}{8} + \frac{2}{3}} \end{aligned}$$



7. If the cylindrical coords of a point are $(2\sqrt{3}, 3\pi/4, 2)$ find the spherical coords of the point.

$$(r, \theta, z) = (2\sqrt{3}, \underline{3\pi/4}, 2) \rightarrow (\rho, \underline{\theta}, \varphi) = \boxed{(4, 3\pi/4, \pi/3)}$$

$$x = 2\sqrt{3} \cos \frac{3\pi}{4} = -\sqrt{6}$$

$$y = 2\sqrt{3} \sin \frac{3\pi}{4} = \sqrt{6}$$

$$z = 2$$

$$\rho^2 = (-\sqrt{6})^2 + (\sqrt{6})^2 + (2)^2$$

$$\rho^2 = 16 \Rightarrow \rho = 4$$

$$2 = 4 \cos \varphi$$

$$\cos \varphi = \frac{1}{2} \Rightarrow \varphi = \frac{\pi}{3}$$